

PACKAGE LEAFLET

CARBOSYLANE CHILD capsule

Read all of this leaflet carefully before you start taking this medicine. It contains important information on your treatment

If you have any further questions, if you have any doubt, ask your doctor or pharmacist

Keep this leaflet, you may need to read it again.

If you need further information and advices, please ask your pharmacist.

If symptoms worsen or persist, consult your doctor.

If you notice any side effects not listed in this leaflet or if any of the side effects gets serious, please tell your doctor or pharmacist.

In this leaflet:

1. WHAT CARBOSYLANE CHILD, capsule IS AND WHEN IT IS USED FOR?
2. WHAT IS THE INFORMATION TO BE AWARE OF BEFORE USING CARBOSYLANE CHILD, capsule?
3. HOW TO USE CARBOSYLANE CHILD, capsule?
4. WHAT ARE THE POSSIBLE SIDE EFFECTS?
5. HOW TO STORE CARBOSYLANE CHILD, capsule?
6. FURTHER INFORMATION

1. WHAT CARBOSYLANE CHILD, capsule IS AND WHEN IT IS USED FOR?

ANTIPLATULENT/ INTESTINAL ADSORBENT

This medicine is indicated in children over 6 years of age for the symptomatic treatment of dyspeptic states (indigestion) and meteorism (intestinal distension).

2. WHAT IS THE INFORMATION TO BE AWARE OF BEFORE USING CARBOSYLANE CHILD, capsule ?

Never use CARBOSYLANE CHILD, capsule in the following cases:

- Allergy to any of the constituents of CARBOSYLANE CHILD, capsule
- In children under 6 years of age (risk of choking of the capsule).
- Bowel obstruction

IF IN DOUBT, IT IS ESSENTIAL TO ASK YOUR DOCTOR OR PHARMACIST FOR ADVICE.

Take special care with CARBOSYLANE CHILD, capsule:

If your child takes digitalis-type drugs, allow an interval between taking CARBOSYLANE and digitalis-type drugs (more than 2 hours, if possible).

Taking or using other medicines

This medicine can reduce the uptake of other medicines, and these should therefore be administered at an interval from it.

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Pregnancy and breast feeding

This medicine, under normal conditions of use, can be used occasionally during pregnancy and breast feeding.

Generally, ask your doctor or pharmacist for advice before taking any medicine.

3. HOW TO USE CARBOSYLANE CHILD, capsule ?

Posology

FOR CHILDREN OVER 6 YEARS OF AGE ONLY.

This medicine is presented in the form of a blue capsule (whose active ingredients are released in the stomach) and a red capsule (not degraded in the stomach)

The usual dosage is

- 2 doses daily for children aged 6 to 10 years
- 3 doses daily for children aged 10 to 15 years

Method of administration

Oral route.

Frequency of administration

The 2 capsules are to be administered at the same time, preferably at the start of the meal.

If you use more CARBOSYLANE CHILD, capsule, than you should:

No case of overdose has been reported, however use of high dose levels can result in dark colouration of the stools.

4. WHAT ARE THE POSSIBLE SIDE EFFECTS ?

Like all medicines, CARBOSYLANE CHILD, capsule can cause side effects, although not everybody gets them.

- dark discolouration of the stools may occur.

Statement of the side effects:

If you feel any adverse effect, tell it to your doctor or your pharmacist or your nurse. This also applies to any adverse effect which would not be mentioned in this leaflet. You can also declare adverse effects directly via the national system of statement: the National Agency for safety of medicines and health products (Ansm) and the Regional Centers of Pharmacovigilance. The web site: www.ansm.sante.fr. By reporting the adverse events, you contribute to supply more information on the medicine safety.

5. HOW TO STORE CARBOSYLANE CHILD, capsule?

Keep out of the reach and sight of children.

Do not use CARBOSYLANE CHILD, capsule, after the expiry date which is stated on the outer packaging.

No special storage conditions.

Medicines should not be disposed of via waste-water or household wastes. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

6. FURTHER INFORMATION

What does CARBOSYLANE CHILD, capsule, contain?

The active substances are:

Gastro-soluble blue capsule

Activated charcoal.....	70.0 mg
Simeticone.....	22.5 mg

For one capsule N° 4 of 107.5 mg.

Gastro-resistant red capsule

Activated charcoal.....	70.0 mg
Simeticone.....	22.5 mg

For one capsule N° 4 of 107.5 mg.

The other ingredients are:

Gastro-soluble blue capsule

Sorbitan oleate, polysorbate 80

Composition of the capsule shell:

Gelatin, indigotin (E132), sulphur dioxide, titanium dioxide (E171).

Gastro-resistant red capsule

Sorbitan oleate, polysorbate 80

Composition of the capsule shell:

Gelatin, erythrosine (E127), titanium dioxide (E171), sulphur dioxide, indigotin (E132).

Composition of the gastro-resistant coating :

Cellulose phthalate acetate, diethyl phthalate, methanol, methylene chloride.

What is CARBOSYLANE CHILD, capsule and contents of the outer pack ?

This medicine contains capsules. Box of 48 (24 doses), 96 (48 doses).

Marketing Authorisation Holder/Operating company:

LABORATOIRES GRIMBERG - 19 rue Poliveau - 75005 Paris – France

Manufacturer:

Laboratoires GRIMBERG - ZA des Boutries - 5, rue Vermont - 78700 Conflans Sainte Honorine France

This leaflet was last approved on 27th September 2012.

