

PACKAGE LEAFLET

SOLACY ADULT, capsule

L-cystine, Precipitated sulphur, Vitamin A, Saccharomyces cerevisiae yeast

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get one of side effects, talk to your doctor or pharmacist. This includes any possible side effect not listed in this leaflet. See section 4.
- You must talk to your doctor if you do not feel better or if you feel worse.

Keep out of the reach of children

In this leaflet:

1. WHAT SOLACY ADULT, capsule IS AND WHEN IT IS USED FOR
2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE SOLACY ADULT, capsule
3. HOW TO TAKE SOLACY ADULT, capsule
4. WHAT ARE POSSIBLE SIDE EFFECTS
5. HOW TO STORE SOLACY ADULT, capsule
6. CONTENTS OF THE PACK AND OTHER INFORMATION

1. WHAT IS SOLACY ADULT, capsule AND WHEN IS IT USED

Pharmacotherapeutic group: Cold and cough medication (respiratory system) - ATC code R05X

Symptomatic adjunctive treatment of rhinopharyngeal disorders in adult.

You must talk to your doctor if you do not feel better or if you feel worse.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE SOLACY ADULT, capsule

Never take SOLACY ADULT, capsule:

- if you are allergic (hypersensitive) to the actives substances or one of the other ingredients of contained in this medicine listed in section 6

Warnings and precautions

Talk to your doctor or pharmacist before taking SOLACY ADULT, capsule.

This medicine contains vitamin A; other medicines contain it. Do not combine them in order not to exceed the maximum recommended dose (see the section on dosage).

Due to the presence of sucrose, this medicine is contra-indicated in patients with fructose intolerance (rare metabolic diseases).

You should not take SOLACY ADULT, capsule if you take a medicine containing retinoids (derivates of vitamin A) due to a risk suggesting hypervitaminosis A.

You should not take an antibiotic of the class of cyclines associated to SOLACY ADULT, capsule at the dosage providing 10 000 IU/day or more of vitamin A because of the risk of intracranial hypertension.

Other medicines and SOLACY ADULT, capsule

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Pregnancy and breast feeding

For pregnant or breastfeeding women, given the daily dietary intake, it is recommended not to exceed a daily dose of 5000 IU of vitamin A brought by the medicines.

As a result, do not exceed the dosage recommended by your doctor.

If you are pregnant or breast-feeding, if you think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

SOLACY ADULT, capsule contains sucrose

3. HOW TO TAKE SOLACY ADULT, capsule

Posology

FOR ADULT USE ONLY.

The recommended dose is 3 capsules daily for 3 months.

Method and route of administration

Oral route.

Swallow the capsules with a glass of water, preferably during a meal.

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist in case of doubt.

If you have taken more SOLACY ADULT, capsule, than you should have:

In case of erroneous intake of an excessive dose, signs of vitamin A overdose may occur. Consult immediately your doctor or your pharmacist.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them. Some cutaneous reactions have been reported.

Reporting of side effects:

If you get any side effect, talk to your doctor or your pharmacist or your nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system: National Agency for safety of medicines and health products (ANSM) and network of the Regional Centers of Pharmacovigilance. Web site: www.signalement-sante.gouv.fr.

By reporting side effects, you can help provide more information on the safety of this medicine.

5. HOW TO STORE SOLACY ADULT, capsule

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the cardboard box after EXP. The expiry date refers to the last day of that month.

Store at a temperature not exceeding 30°C.

Do not throw away any medicine via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What does SOLACY ADULT, capsule, contain?

- The active substances are:

L-Cystine.....	72.6 mg
Precipitated sulphur.....	22.0 mg
Coated vitamin A acetate quantity equivalent to retinol.....	1650 IU
Saccharomyces cerevisiae yeast.....	77.4 mg

Per a capsule

- The other ingredients are:
Magnesium stearate, gelatin, titanium dioxide (E 171), antioxidant (BHT), sucrose, maize starch.

What SOLACY ADULT, capsule looks like and contents of the pack?

This medicine contains capsules. Box of 45 and 90 capsules.

Marketing Authorisation Holder:

Laboratoires GRIMBERG SA – 44 avenue Georges Pompidou- 92300 Levallois-Perret - France

Manufacturer:

Laboratoires GRIMBERG SA - ZA des Boutries - rue Vermont - 78704 Conflans Sainte Honorine cedex - France

This leaflet was last revised in January 2019.

Detailed information on this medicinal product is available on ANSM website (France)